

### ALICE 3




复旦大学 (上海,中国) 2025-11-09

Jochen Klein (CERN)


### Frustration

- Many reasons to believe in formation of deconfined state of matter in ultra-relativistic heavy-ion collisions, BUT:
  - no single measurement to unequivocally demonstrate deconfinement
  - no direct measurement of the initial conditions, e.g. temperature
  - no direct demonstration of phase transition
  - missing handles on interaction potentials and nature of bound states



### Deconfinement


- What do we need to demonstrate deconfinement?
  - traceable probe embedded in quark-gluon plasma
    - → light quarks and gluons produced thermally → not traceable 👎
    - → heavy quarks only produced initially → traceable 👍
    - → charm quarks equilibrate → embedded in QGP 👍
  - proof of free motion within quark-gluon plasma
    - → charm quarks produced independently and well separated
    - → combination of multiple charm quarks demonstrates motion de
  - exclusion of other production mechanisms
    - → formation of multi-charm states from fragmentation strongly suppressed 👍
- Multi-charm hadrons are ideal probe to unequivocally demonstrate equilibration and deconfinement



unique chance at LHC, but not possible yet

### Initial conditions


- What do we need to measure initial temperature?
  - hadrons produced at freeze-out temperature
    - → information about initial state mostly lost 👎
  - electromagnetic probes produced throughout the evolution
    - → no strong interaction → escape from quark-gluon plasma de
  - photons suffer from blue shift
  - di-leptons escape and invariant mass is unaffected by blue shift description.
- Di-leptons are ideal probes to measure properties of the early phases



needs to be done at LHC but not possible yet

### Interaction potentials

- What do we need to understand interaction potentials and the nature of bound states?
  - production yields only give indirect insights
  - correlations of compounds carry information
     on interaction potentials and possible bound states
- Momentum correlations are ideal probes
   to unravel interaction potentials between hadrons and nature of bound states




r<sub>molecule</sub>
as large as 5 fm

unique chance at LHC, but not possible yet

### Phase transition

- What do we need to detect and characterise the phase transition of QCD?
  - phase transition characterised by critical fluctuations
    - → cumulants of conserved quantities
    - → net-baryon cumulants can be measured
    - → can be confronted with predictions from lattice QCD<sub>50</sub>
- Net-baryon fluctuations are ideal probes to probe the nature of the phase transition



unique chance at LHC, but not possible yet

### Key measurements

#### • (Multi-)heavy-flavoured probes


- modified parton shower
- \*\*\* transport properties
- hadronisation

#### Dielectrons down to low mass

- temperature and early stage
- chiral symmetry restoration

#### Correlations and fluctuations

- net-baryon fluctuations
- \*\*\* transport properties



electron identification

Background from heavy-flavour decays  $c\bar{c} \rightarrow D\bar{D} \rightarrow e^+ e^- ...$ 

#### **Experimental requirements**

- Excellent pointing resolution
- Tracking down to  $p_T \approx 0$
- Excellent particle identification
- Large acceptance
- High rates for large data samples


Progress relies on detector performance and statistics


### Observables

- Heavy-flavour hadrons (p<sub>T</sub>  $\rightarrow$  0,  $|\eta|$  < 4)
  - vertexing (decay chain)
  - tracking (inv. mass resolution)
  - hadron ID (background suppression)
- **Dielectrons** (p<sub>T</sub> ~0.1 3 GeV/c,  $M_{ee}$  ~0.1 4 GeV/ $c^2$ )
  - vertexing (HF background suppression)
  - tracking (inv. mass resolution)
  - electron ID
- **Photons** (100 MeV/c 50 GeV/c, wide  $\eta$  range)
  - electromagnetic calorimetry
- Quarkonia and Exotica ( $p_T \rightarrow 0$ )
  - muon ID
- Ultrasoft photons ( $p_T = 1 50 \text{ MeV/}c$ )
  - dedicated forward detector
- Nuclei
  - identification of z > 1 particles

#### **Key requirements**

- Tracking over large rapidity range
- Excellent vertexing
- Excellent particle identification
- High rate







### Detector concepts


How do we optimise a detector for this programme?

## Motion of charged particles

 Motion of charged particle in a magnetic field determined by Lorentz force

$$\overrightarrow{F} = q\left(\overrightarrow{v} \times \overrightarrow{B}\right) \Rightarrow \dot{\overrightarrow{v}} = \frac{q}{\gamma m} \left(\overrightarrow{v} \times \overrightarrow{B}\right)$$

- force perpendicular to direction of motion
  - in magnetic field only change of direction
  - → no change of energy → γ constant
- valid even relativistically (despite non-covariant formulation)



### Helix motion in homogeneous field

- Charged particle in homogeneous magnetic field
  - → helix motion coaxial with magnetic field (in z direction)

$$x = \frac{v_{\mathrm{T}}}{\eta \omega_{B}} \sin(\eta \omega_{B} t + \psi_{0}) + x_{0} \qquad y = \frac{v_{\mathrm{T}}}{\eta \omega_{B}} \cos(\eta \omega_{B} t + \psi_{0}) + y_{0} \qquad z = v_{3} t + z_{0}$$
with cyclotron frequency  $\omega_{B} = \frac{|q|B}{\gamma m}$  and  $\eta = q/|q|$ 

• radius of curvature given by  $p_{\perp} = |q|BR$ , when using standard units (GeV/c, T, m)  $\rightarrow p_{\perp} = 0.3 BR$ 


1 GeV/c particle in 1 T field

→ ~3 m radius

curvature in magnetic field directly linked to (transverse) momentum


## Magnetic field configurations


- Dipole fields
  - often used in fixed-target experiments or forward part of collider experiments
- Solenoidal fields
  - often used in collider experiments (rotationally symmetric)



- Toroidal fields
  - often used as extension of solenoidal fields (perpendicular, rotationally symmetric)


Deflection depends on motion perpendicular to magnetic fields





## Tracking

- Reconstruction of trajectory of a particle through the detector with the goal of determining properties of the particle, in particular momentum
  - connection of hits in one or more detectors (originating from the same particle)
    - **"→ track finding**
  - fitting of track parameters from hits (taking into account interaction with material and magnetic fields)
    - **"→** track fitting
- Requirements for tracking detectors
  - 3 space points fully describe trajectory (under ideal conditions)
  - additional measurements can reduce uncertainties
  - need for redundancy and suppression of fake tracks (occupancy)



## Criteria for tracking performance

#### Ideal reconstruction

- assignment of all hits to correct track
  - → all tracks are reconstructed, no impact from wrongly assigned hits
- precise estimation of track parameters
  - → unbiased, minimal uncertainties

#### • Figures of merit


- acceptance → fraction of area covered by detector
- efficiency → fraction of reconstructed tracks
- fake hit probability → probability of assigning hits from other particles
- extrapolation uncertainties → propagation beyond volume covered by tracker
- momentum resolution

## Sagitta

- Momentum measurements boils down to
   measurement of curvature in magnetic field
   → extraction from bubble chamber images
- Sagitta → maximum deviation
   from line between endpoints of track segment

$$s = R - R \cos \frac{\vartheta}{2} = 2R \sin^2 \frac{\vartheta}{4} \approx \frac{R\vartheta^2}{8} \approx \frac{L^2}{8R} = \frac{qBL^2}{8p_{\perp}}$$

- quadratic dependence on lever arm
- relative uncertainty of momentum proportional to relative uncertainty of sagitta




geometric approach, next consider measurement of points along trajectory

### Position resolution

- Measurement of hit position limited by detector resolution
  - uncertainty for rectangular structure with binary readout pitch p

$$\sigma_{\text{pos}}^2 = \int_{-p/2}^{p/2} dx \, x^2 \frac{1}{p} = \frac{p^2}{12} \Rightarrow \sigma_{\text{pos}} = \frac{p}{\sqrt{12}}$$

- uncertainty reduced when signal spreads across multiple bins (charge sharing), possibly with information on charge per bin
- gas detectors typically  $\mathcal{O}(100~\mu m)$ , semiconductor detectors typically  $\mathcal{O}(10~\mu m)$



## Multiple scattering

- Material (incl. sensitive volume!) leads to multiple scattering
  - → random change of direction on top of curvature in magnetic field
  - width of distribution

$$\sigma_{\alpha} = \frac{0.0136 \,\text{GeV/}c}{\beta p} \sqrt{\frac{d}{X_0}}$$

- inversely proportional to momentum
- scales with square root of material thickness
- Multiple scattering poses fundamental limit on measurement precision
  - → cannot be mitigated by improved position resolution
  - angular effect, i.e.  $\sigma_x \propto \sigma_\alpha \cdot \Delta x$

### Track fit

- Global  $x^2$  minimisation of parameterised function f(x) for measured points  $(x_i, y_i)$ 
  - assume linear dependency on parameters, e.g.
    - straight line

$$\rightarrow y_i = f(x_i) = a_0 + a_1 x_i$$

• parabola (as approximation to circular shape)

$$\rightarrow$$
 y<sub>i</sub> = f(x<sub>i</sub>) = a<sub>0</sub> + a<sub>1</sub> x<sub>i</sub> + a<sub>2</sub> x<sub>i</sub><sup>2</sup>/2 (a<sub>2</sub> = 1/R)

• find parameters a to minimise

$$\chi^2 = (y - Ga)^T W (y - Ga)$$
 with  $G := x_i^j$  and weights  $W$  (W is identity matrix for equal weights)

• minimum achieved for **solution of normal equation**  $G^TWGa = G^TWy$ solvable by matrix inversion or as system of linear equations (numerically more stable)

## Tracking uncertainties

- Uncertainties of measurements described by **covariance matrix**  $C_v = E((y Ey)(y Ey)^T)$
- Propagation of uncertainties to parameters a  $C_a = BC_vB^T$  with  $B = (G^TWG)^{-1}G^TW$  (from previous slide)
- Optimal choice of W to achieve unbiased parameter estimation with minimal uncertainties  $W=C_{\rm v}^{-1}$
- Uncertainties for ideal weights are  $C_a = (G^T C_v^{-1} G)^{-1}$

#### Momentum resolution

- Transverse momentum extracted from curvature in magnetic field:  $p_{\perp} = |q| BR$ 
  - approximated circle by parabola  $f(x_i) = a_0 + a_1 x_i + a_2 x_i^2 / 2$  with  $a_2 = 1/R$
  - momentum resolution proportional to uncertainty of curvature:

$$\frac{\Delta p_{\mathrm{T}}}{p_{T}} = \frac{\Delta R}{R} = \frac{\Delta a_{2}}{a_{2}} = \frac{p_{\mathrm{T}}}{|q|B} \Delta a_{2}$$

Position resolution of N+1 equidistant layers leads to uncertainty

$$\frac{\Delta p_{\rm T}}{p_{\rm T}} = \frac{\sigma p_{\rm T}}{|q|BL^2} \sqrt{\frac{720N^3}{(N-1)(N+1)(N+2)(N+3)}} \approx \frac{\sigma p_{\rm T}}{|q|BL^2} \sqrt{\frac{720}{N+4}}$$
(Glückstern formula, NIM 24 (1963) 381)

Multiple scattering in N+1 equidistant layers leads to uncertainty

$$\frac{\Delta p_{\rm T}}{p_{\rm T}} = \frac{N}{\sqrt{(N+1)(N-1)}} \frac{p\sigma_{\alpha}\sqrt{N+1}}{BL} = \frac{N}{\sqrt{(N+1)(N-1)}} \frac{0.0136\,{\rm GeV}/c\,\sqrt{d_{\rm tot}/X_0}}{\beta BL}$$

## Considerations for detector layout


• Multiple scattering contributes

$$\propto \frac{\sqrt{d_{\text{tot}}/X_0} \cosh \eta}{\beta BL}$$

- independent of momentum (for  $\beta \approx 1$ )
- linear dependence on lever arm
- scales with square root of material
- Position resolution contributes

$$\propto \frac{\sigma p_{\mathrm{T}}}{BL^2}$$

- linear rise with momentum
- quadratic dependence on lever arm




m.s. sets irreducible limit, p.r. takes over at large momenta


## Track finding

- So far track fit assuming knowledge of contributing hits
  - → in reality need to identify hit-to-track association from data
  - → task of pattern recognition
- Variety of methods available (and used), usability often determined by computational effort
  - fit all combinations (and reject based on  $\chi^2$ )
  - Hough transformation
  - cellular automaton
  - Kalman filter
  - machine learning

## Hough transformation

- Idea: transformation from space coordinates to parameter coordinates,
   e.g. describing a straight line y = ax + b
  - every hit (x, y) transforms into a line in the parameter space
  - parameter lines intersect in a single point (ideal case) or cluster around a value (with uncertainties)
  - accumulation point in the parameter space transforms into a line in real space
- Can be generalised to more complex parameterisations





### Cellular automaton

- Connect hits from adjacent layers within a search window
  - → tree of connections
  - combine track segments
  - longest paths are candidates
  - select candidates based on track fit
- Advantages
  - local matching of hits
    - → contain computational effort





Figure by M. Puccio

### Kalman filter

- Idea: propagate track through detector and update parameters
  - → combination of track finding and fitting steps
  - Define parametrisation of track at given point together with propagator to next layer
    - propagator can take into account effects such as energy loss in material
  - Find a seed, i.e. rough track parametrisation from a single hit or track segment
  - Iterate over layers
    - propagate track to next layer
    - update parameterisation based on hit in this layer
- Algorithm can be repeated in opposite direction for refined fit and smoothing



## Detector optimisation


- Impact of number of layers on
  - track finding
  - track fitting
- Impact of placement of layers on
  - track finding
  - track fitting
- Impact of lever arm and magnetic field on
  - momentum resolution
- Impact of dead zone and inefficiencies on
  - corrections for analysis

### Momentum resolution


• Consider momentum resolution in solenoidal field

• 
$$\propto \frac{\sqrt{d_{\mathrm{tot}}/X_0}\,\cosh\eta}{\beta BL}$$
 for multiple scattering

- $\propto \frac{\sigma p_{\rm T}}{BL^2}$  for position resolution
- Objectives
  - choose **lever arm L** required for momentum resolution: area (and cost) scales quadratically with L (for fixed  $\eta$  coverage)
  - choose magnetic field:
     higher field improves mom. resolution, limits acceptance, increases magnet cost
  - optimise **number of layers**: more layers add material, help with track finding, increase cost
  - minimise material per layer: challenge on power consumption, cooling, mechanics



Trajectory through cylindrical layers



forward coverage with barrel layers

- large areas → expensive
- deterioration of performance

## Tracker layout

- Objectives
  - optimise momentum resolution
  - reduce instrumented area
  - minimise path length in material (avoid shallow angles)
- Considerations
  - Barrel layers well suited to cover central region (up to  $\eta \approx 1.5$ )
  - **Disks** well suited to cover forward region (beyond  $\eta \approx 1.5$ )
  - Inclined layers can be used for transition region
- Ideal layout depends on relative contributions of
  - multiple scattering
  - position resolution
  - track finding (pattern recognition)

## Mismatch probability

#### Objective

• minimise assignment of hits not originating from a given track

#### A conceptual approach

 probability of wrong assignment scales with number of hits in vicinity


$$P \propto \frac{\Delta R^2}{R^2} \cdot x/X_0$$

here: assuming quadratic decay of particle flux

distance from preceding layer to achieve constant probability

$$\Delta R \propto \sqrt{\frac{P}{x/X_0}} \cdot R$$

- Inefficiencies and dead areas can lead to significant deterioration of matching
  - layers often arranged in pairs to avoid lack of position information



### Additional considerations

#### Momentum range

- limited by particles not reaching outer layers
- limited by momentum resolution

#### Propagation to other detectors

limited by pointing resolution towards other (outer) detectors

#### • Track length determination

• limited by precision of propagation inside tracker


#### Reconstruction of secondary particles


 limited by number of hits for particles produced at a distance from the primary vertex


ALICE 3 | 2025-11-09 | Jochen Klein

#### Vertices

- Reconstruction of primary vertex,
   i.e. point of underlying interaction,
   from emerging tracks
  - need to distinguish primary vertices
  - assignment of tracks to interactions
- Reconstruction of decay vertices
  - distinction of prompt and non-prompt particles
  - reconstruction of decay vertices and decay chains
- Need for pointing resolutions well below 100 μm









## Track propagation

- Track finding like for tracking
- Track and propagation defined by straight line through 2 hits
  - magnetic field can often be neglected because of short distances
  - typically vertex detectors feature at least three layers for track finding
- Extrapolation limited by
  - position resolution
  - multiple scattering







### Position resolution

- Impact parameter obtained from track extrapolation
  - approximation by straight line  $f(x_i) = a_0 + a_1 x_i$
  - with first layer at radius  $r_0$ , uncertainty given by  $\sigma_{xy} = \sigma_{a_1} \cdot r_0$   $\sigma_z = \sigma_{a_1} \cdot r_0 \cdot \cosh \eta$



- $\sigma_{d_{xy}} \propto \sigma_{r\varphi}$  (transverse impact parameter)
- $\sigma_{d_z} \propto \sigma_z$  (longitudinal impact parameter)






## Multiple scattering

Multiple scattering in N+1 equidistant layers leads to

$$\sigma_{\varphi} \propto \frac{\sqrt{d/X_0 \cosh \eta}}{\beta p_{\mathrm{T}}}$$


$$\sigma_{\vartheta} \propto \frac{\sqrt{d/X_0 \cosh \eta}}{\beta p} \propto \frac{\sqrt{d/X_0}}{\beta p_{\mathrm{T}} \sqrt{\cosh \eta}}$$



Propagation towards interaction point leads to

• 
$$\sigma_{d_{xy}} \propto \sigma_{\varphi} \cdot r_0 \propto \frac{\sqrt{d/X_0 \cosh \eta}}{\beta p_{\mathrm{T}}} r_0$$
 (transverse)


• 
$$\sigma_{d_z} \propto \sigma_{\vartheta} \cdot r_0 \cdot \cosh^2 \eta \propto \frac{\sqrt{d/X_0}}{\beta p_{\mathrm{T}}} \cdot r_0 \cdot \sqrt{\cosh \eta}^3$$
 (longitudinal)



## Considerations for detector layout

#### Pointing resolution scales

- with distance of closest layer
  - → priority to be as close as possible
- with square root of material (m.s.  $\propto 1/p$ )
  - → minimise material
- with position resolution
  - → keep sub-dominant (within range)
- More layers do not improve precision but can be needed for track finding



ALICE 3 | 2025-11-09 | Jochen Klein

# Scaling of resolutions (for cylindrical layers)

position resolution

multiple scattering

momentum

$$\propto \frac{\sigma p_{\rm T}}{BL^2}$$

transverse DCA

$$\propto \sigma_{xy}$$

**longitudinal DCA** 

$$\propto \sigma_{z}$$

$$\propto \sqrt{d_{\text{tot}}/X_0} \frac{\sqrt{\cosh \eta}}{\beta BL}$$

$$\propto r_0 \sqrt{d/X_0} \frac{\sqrt{\cosh \eta}}{\beta p_{\text{T}}}$$

$$\propto r_0 \sqrt{d/X_0} \frac{\sqrt{\cosh \eta}}{\beta p_{\text{T}}}$$

$$\propto r_0 \sqrt{d/X_0} \frac{\sqrt{\cosh \eta}}{\beta p_{\text{T}}}$$

#### Particle identification


- Mass measurement from combination of tracking and additional measurements
  - Specific energy loss
    - deposited charge (time over threshold)
  - Time of flight separation  $\propto L/\sigma_{\rm tof}$ 
    - large path length with fast time resolution
  - Cherenkov radiation with angle  $\cos \vartheta = 1 / n\beta$ 
    - refractive index to optimise coverage
    - → angular resolution
  - Electromagnetic shower to identify electrons
  - Absorption to identify muons

Combination of techniques to achieve PID goals

## Bethe-Bloch equation

$$-\frac{\mathrm{d}E}{\mathrm{d}x} = Kz^2 \frac{Z}{A} \rho \frac{1}{\beta^2} \left[ \frac{1}{2} \log \frac{2m_e c^2 \beta^2 \gamma^2 T_{\text{max}}}{I^2} - \beta^2 - \frac{\delta}{2} \right]$$


- Calculation of specific energy loss by Bethe and Bloch
  - T<sub>max</sub>: max. energy transfer (in single collision)
  - I: ionisation potential ( $\sim$ (10 ± 1) Z eV for elements beyond O)
  - δ/2: density correction (Lorentz contraction + polarisability of material)



# Specific energy loss

- Deposited energy in any detector is function of  $\beta \gamma = p/m$ 
  - for given momentum
    - → function of mass
  - ambiguities at line crossings
- Requires combined measurement of
  - curvature → momentum
  - deposited energy → dE/dx
- Well suited for integration in tracker with readout of deposited energy

#### **ALICE TPC**




# Time of flight

- Velocity of particle (at given momentum) depends on mass
  - → different time of flight for different mass hypotheses:

$$\Delta t = \frac{L}{c} \left( \frac{1}{\beta_1} - \frac{1}{\beta_2} \right) \approx \frac{Lc}{2p^2} (m_1^2 - m_2^2)$$

- Separation power  $N_{\sigma} = \frac{\mid m_1^2 m_2^2 \mid L}{2p^2 \sigma_t c}$ 
  - improves with
  - path length
  - time resolution → need for fast detectors,
     e.g. scintillators, Cherenkov, MRPCs, LGADs
  - momentum resolution



40

ALI-PERF-106336


- Often realised as combination of fast detector outside of tracker
  - → no impact on tracking performance, large path length

# 4d tracking

- Time information can be used as additional coordinate in track finding and fitting
  - reduce mismatch probability in high occupancy environment, e.g. from pile-up, i.e. multiple collisions in short sequence
  - required time resolution depends on nature of pile-up
  - integration of time measurement in
    - every tracking layer
      - → so far only feasible with moderate time resolution (material)
    - dedicated timing layers
      - → reduce tracks to be considered for each collision

### Cherenkov detectors

- Cherenkov effect
  - → emission of light by particles above speed of light
  - presence of radiation → threshold Cherenkov
  - emission angle function of  $\beta = p/E$ 
    - $\rightarrow$  combination of  $\beta$  and p give access to mass
- Measurement of Cherenkov angle requires
  - sufficient production of light
    - → minimum amount of material
  - focusing onto (single-)photon-sensitive sensors
    - → expansion gap or optics
- Typically realised as additional detector outside of tracker (need for space and material)



# Calorimetry

- Detection of total energy based on complete conversion into secondary particles
  - ECal: pair production and bremsstrahlung
  - HCal: nuclear reactions producing pions, ...
- Calorimeters are destructive detectors
  - → placed outside of trackers
- Propagation of (charged particle) tracks to calorimeters can be required for additional information, e.g. avoidance of double counting

#### Muon identification


- Exploit unique signature of muons
  - energy loss through ionisation/excitation (minimum ionising particle, Bethe Bloch)
  - no electromagnetic shower (too heavy)
  - no hadronic shower (no strong interaction)
- Absorber can be used to block anything but muons
  - → outermost part of a detector
- Tracking can be needed for
  - matching to tracks before absorber
  - propagation of tracks through absorber

## Summary part 1

- Fundamental physics questions remain open
  - can be addressed at the LHC, but not with existing experiments
- Key measurements need to drive optimisation of detector
  - tracking
  - vertexing
  - particle identification

Time for discussion ©







# Backup


## Quantum Chromodynamics

Lagrangian of **Quantum Chromodynamics** → strong interaction

$$\mathcal{L}_{\text{QCD}} = \overline{\psi}_i \left( i \gamma^{\mu} \left( D_{\mu} \right)_{ij} - m \delta_{ij} \right) \psi_j - \frac{1}{4} G^a_{\mu\nu} G^{\mu\nu}_a$$

- Interactions with high Q<sup>2</sup> well described by perturbative calculations
- Many non-trivial features of QCD emerge from low-Q2 regime
  - confinement → no free quarks
  - chiral symmetry breaking → nucleon mass
  - fragmentation → formation of hadrons
  - QCD matter → thermodynamic properties


**Emerging features of QCD of fundamental interest** 




## QCD matter

- Nuclear matter → QCD matter at ambient conditions
- ullet QCD matter at different temperatures T and densities  $\mu_B$ 
  - deconfinement for  $T \to \infty$ ,  $\mu_B \to \infty$
  - chiral symmetry restoration
  - superconductivity
- Numerical calculations of QCD on a discretised lattice
  - **cross-over** from hadron gas to quark-gluon plasma around pseudo-critical temperature T<sub>pc</sub>
  - interactions still relevant for  $T \gg T_{pc}$


Experimental data crucial to understand QCD matter





#### Nuclear collisions

- Ultra-relativistic collisions of nuclei
  - → hot and dense nuclear matter
  - deconfinement of nuclear matter to quark-gluon plasma and back
  - small, short-lived, dynamic system
  - self-generated probes from all stages of the evolution
  - conditions controlled through
     collision energy and geometry



Electromagnetic radiation (  $\propto T^2$  )

Hadron momentum distributions, azimuthal anisotropy

Hadron abundances 'hadrochemistry'


**Hadron correlations, fluctuations** 


Ultra-relativistic heavy-ion collisions

→ excellent means to study hot QCD matter

# Quark-gluon plasma

- Consider QCD matter in heavy-ion collision
  - → path in phase diagram
  - kinetic energy → creation of quark-gluon plasma
  - expansion and cool down → hadron production
- What is the initial temperature and how does it evolve?
  - → thermal spectra of photons and dileptons
- What are the mechanisms for the restoration of chiral symmetry in the quark-gluon plasma?
  - $\rightarrow \rho$ -a<sub>1</sub> mixing  $\rightarrow$  dileptons
- What is the nature of the phase transition?
  - → critical behaviour → fluctuations (cf. lattice QCD)








104

## Interactions with QGP

- Consider QGP phase of a heavy-ion collision
  - quarks and gluons interact with the plasma
  - pressure-driven expansion
    - → viscous, relativistic hydrodynamics
- What is the nature of interactions between quarks and gluons and the quark-gluon plasma?
  - mass dependence of medium interaction
    - → charm and beauty jets
  - transport coefficients for charm and beauty
    - $\rightarrow$  anisotropy (v<sub>n</sub>) and suppression (R<sub>AA</sub>)
  - equilibration of charm and beauty
    - → hadron yields and decorrelation







### Hadron formation

- Consider transition from QGP to hadrons
  - hadron yields described by thermal models with temperature and baryochemical potential
  - light-flavour quarks produced thermally, heavy-flavour quarks only in initial scatterings
- To what extent do quarks of different mass reach thermal equilibrium?
  - statistical hadronisation of charm/beauty quarks
     → yields of heavy-flavour hadrons
- How do quarks and gluons transition to hadrons as the quark-gluon plasma cools down?
  - combination of independently produced quarks
     → multi-heavy-flavour states
  - production of bound states
    - → hyper/charm-nuclei

