



復旦大學
FUDAN UNIVERSITY

Quantum Entanglement Studies
in Nuclear and Particle Physics



University of
Pittsburgh®

Quantum information in DIS process at the EIC

Based on arXiv:2510.23773, PRD(2026), **Kun Cheng**, Tao Han and Sokratis Trifinopoulos

Rich Quantum Information at colliders

- ▶ LHC, Belle, BESE, RHIC...
 - ▶ $t\bar{t}, b\bar{b}, W^\pm, Z, q\bar{q}, \Lambda\bar{\Lambda}, \tau^\pm \dots$,
-
- A near future collider — Electron Ion Collider (EIC).
 - ▶ Connections with QCD (nucleon tomography)
 - ▶ More control on the initial states (polarization, ...)
 - ▶ Is the special properties of the EIC relevant to quantum information?

Spin states at colliders

- Depends on initial state, interactions, and kinematics
 - ▶ The “unnormalized” spin state of final state $\mathcal{AB}$

$$|\mathcal{AB}\rangle = \hat{\mathcal{T}} |\text{initial}\rangle$$


$$\hat{R}_{\mathcal{AB}}(\mathbf{k}) = \mathcal{M}_{\alpha\bar{\alpha}}^{\lambda}(\mathbf{k}) \rho_{\lambda\lambda'}^{\text{initial}} (\mathcal{M}_{\alpha'\bar{\alpha}'}^{\lambda'}(\mathbf{k}))^* |\alpha\bar{\alpha}\rangle \langle\alpha'\bar{\alpha}'|$$

If the initial state is a superposition/mixture of ① and ②, then final state is a superposition/mixture of the states produced by ① and ②,

- E.g. one of the most studied: $gg \rightarrow t\bar{t}$
- E.g. no entanglement for $q\bar{q} \rightarrow t\bar{t}$ near threshold:
both $q_L\bar{q}_R$ and $q_R\bar{q}_L$ produce separable states, mixture cannot generate entanglement

$$|\psi^{\mathcal{AB}}\rangle = |\psi^{\mathcal{A}}\rangle \otimes |\psi^{\mathcal{B}}\rangle$$

$$\rho^{\mathcal{AB}} = \sum_i p_i \rho_i^{\mathcal{A}} \otimes \rho_i^{\mathcal{B}}$$



[Afik and de Nova, Quantum (2022)]

Quantum information at the EIC

- At the EIC, $g\gamma/\gamma\gamma \rightarrow q\bar{q}$ is a similar process (at lowest order)

[Qi, Guo, and Xiao, PRD (2026)]

[Fucilla and Hatta, PRD (2026)]

[Hatta and Schoenleber, arXiv:2511.04537]

[Fucilla, Hatta, and Xiao, arXiv:2604.11697]



- $gg \rightarrow q\bar{q}$, $g\gamma \rightarrow q\bar{q}$ and $\gamma\gamma \rightarrow q\bar{q}$ produces exactly the same spin state [KC, Han and Low, PRD (2025)]

- Something typical/special

- Virtuality of photon
 - Ultrapерipheral Collisions (UPC)
 - Linear polarization of gluon

- What about the leading process $eq \rightarrow eq$?

[Afik and de Nova, Quantum (2022)]



[Qi, Guo, and Xiao, PRD (2026)]

Leading processes at the EIC

- Serves for many major physic focus of the EIC :
 - ▶ “3D tomography of the nucleon”
 - ▶ PDF, TMD PDF, polarized PDF, tensor charge...

[EIC white paper, arXiv:1212.1701]




FIG. 1: Elastic $e^- p$ (inelastic $e^- q$) scattering

- What's the quantumness of the elastic and deep inelastic scattering process?
(in the presence of beam polarization)
 - ▶ e.g. entanglement (concurrence $\mathcal{C}$)
 - ▶ e.g. nonstabilizerness (magic $\mathcal{M}_2$)

[see yesterday's talk by Prof. Qun Wang, Hao Zhang and Jinniu Hu]

Leading processes at the EIC

- Helicity is unchanged in t -channel process
 - ▶ $|e_{\uparrow}\rangle \otimes |p_{\alpha}\rangle \longrightarrow |e_{\uparrow}\rangle \otimes |p_{\alpha'}\rangle$
 - ▶ $|e_{\downarrow}\rangle \otimes |p_{\alpha}\rangle \longrightarrow |e_{\downarrow}\rangle \otimes |p_{\alpha''}\rangle$
- e.g. backward limit, $|e_{\uparrow}\rangle \rightarrow |\uparrow\downarrow\rangle, |e_{\downarrow}\rangle \rightarrow |\downarrow\uparrow\rangle$
 - ▶ Unpolarized beams, mixture of $|e_{\uparrow}\rangle$ and $|e_{\downarrow}\rangle$
incoherent mixture of $|\uparrow\downarrow\rangle$ and $|\downarrow\uparrow\rangle$
 - ▶ Any non-trivial quantum information?




FIG. 1: Elastic e^-p (inelastic e^-q) scattering

$$\mathcal{M}_{\alpha'\beta'}^{\alpha\beta} \equiv \mathcal{M}(e^-(\alpha)p(\beta) \rightarrow e^-(\alpha')p(\beta'))$$

$$\mathcal{M}_{\uparrow\uparrow}^{\uparrow\uparrow} \approx 2Q_p e^2 \frac{s+t}{t} e^{i\phi},$$

$$\mathcal{M}_{\uparrow\downarrow}^{\uparrow\downarrow} = \mathcal{M}_{\downarrow\uparrow}^{\downarrow\uparrow} \approx 2Q_p e^2 \frac{s}{t},$$

$$\mathcal{M}_{\downarrow\downarrow}^{\downarrow\downarrow} \approx 2Q_p e^2 \frac{s+t}{t} e^{-i\phi},$$

Leading processes at the EIC

- Helicity is unchanged in t -channel process
 - ▶ $|e_{\uparrow}\rangle \otimes |p_{\alpha}\rangle \longrightarrow |e_{\uparrow}\rangle \otimes |p_{\alpha'}\rangle$
 - ▶ $|e_{\downarrow}\rangle \otimes |p_{\alpha}\rangle \longrightarrow |e_{\downarrow}\rangle \otimes |p_{\alpha''}\rangle$
- e.g. backward limit, $|e_{\uparrow}\rangle \rightarrow |\uparrow\downarrow\rangle, |e_{\downarrow}\rangle \rightarrow |\downarrow\uparrow\rangle$
 - ▶ Unpolarized beams, mixture of $|e_{\uparrow}\rangle$ and $|e_{\downarrow}\rangle$
incoherent mixture of $|\uparrow\downarrow\rangle$ and $|\downarrow\uparrow\rangle$
- **Opportunities at the EIC: transverse polarization**
 - ▶ Incoming beam: superposition of $|e_{\uparrow}\rangle$ and $|e_{\downarrow}\rangle$
 - ▶ Final state: entangled $|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle$





FIG. 1: Elastic e^-p (inelastic e^-q) scattering



[KC, Han and Trifinopoulos, arXiv:2510.23773]

Beam polarizations

- Polarizations & Bloch vector $\rho = \frac{I_2 + b_i \sigma_i}{2}, \quad b_i = \langle \sigma_i \rangle$
 - ▶ Pure state $|\vec{b}| = 1$: e.g. $\begin{cases} b_z = 1 & \longrightarrow |\psi\rangle = |\uparrow\rangle \\ b_x = 1 & \longrightarrow |\psi\rangle = (|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2} \end{cases}$
 - ▶ Mixed state $|\vec{b}| < 1$: e.g. $|\vec{b}| = 10\% \longrightarrow \rho = \underbrace{10\%}_{\text{degree of beam polarization}} |\psi\rangle \langle \psi| + 90\% \frac{I_2}{2}$
- Transverse polarization naturally exists
 - ▶ Sokolov-Ternov effect: ideally >90%
Unpolarized $\implies$ Transverse (electron spin becomes opposite to magnetic field)
 - ▶ Transverse $\implies$ Longitudinal
Transverse polarization is a medium stage even we only want longitudinal polarization

Beam polarizations

- Polarizations & Bloch vector

$$\rho = \frac{I_2 + b_i \sigma_i}{2}$$

- ▶ Pure state $|\vec{b}| = 1$: e.g. $\begin{cases} b_z = 1 \longrightarrow \\ b_x = 1 \longrightarrow \end{cases}$
- ▶ Mixed state $|\vec{b}| < 1$: e.g. $|\vec{b}| = 10\% \longrightarrow \rho$

- Transverse polarization naturally exists

- ▶ Sokolov-Ternov effect: ideally >90%

Unpolarized $\implies$ Transverse (electron spin becomes opposite to magnetic field)

- ▶ Transverse $\implies$ Longitudinal


- ▶ >70% at EIC for both beams

[EIC yellow report, arXiv:2103.05419]

More details in conceptual design report

[www.osti.gov/biblio/1765663/]

Illustration figure from HERA




DIS process with transverse polarization

- E.g. purely transversely polarized

$$|ep/eq\rangle \propto \left(\frac{s - Q^2}{s} \right) (e^{i\phi} |\uparrow\uparrow\rangle + e^{-i\phi} |\downarrow\downarrow\rangle) + |\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle$$

- For either elastic or inelastic scattering, the entanglement of final state occurs only when:
 - ▶ Initial beams are transversely polarized
 - ▶ Q^2 is sizable (then mostly DIS process)
- Entanglement increases with:
 - ▶ $Q^2 \sim 1 - \cos \theta$
 - ▶ beam polarization $\vec{b}_\perp$



The dependence of the entanglement on the polarization gives us the opportunity to control the state production by changing the initial state polarization.

Quark polarization: transversity PDF

- Transversity PDF $h_{1,q}(x)$: when proton spin points along transverse direction $+\hat{n}$
 - ▶ probability of finding q polarized along $+\hat{n}$ minus the probability of finding q polarized along $-\hat{n}$
- Together with the unpolarized PDF $f_q(x)$, probability of finding q from the proton, we see the relationship between the spin density matrices of quark and proton:


$$\rho^p = \frac{\mathbb{1}_2 + \vec{b}_\perp^p \cdot \vec{\sigma}_\perp}{2}, \quad \rho^q = \frac{\mathbb{1}_2 + \vec{b}_\perp^q \cdot \vec{\sigma}_\perp}{2}, \quad \vec{b}_\perp^q = \frac{h_{1,q}(x)}{f_q(x)} \vec{b}_\perp^p$$

- Without changing collider settings, we can choose different quark polarizations.

Quark polarization: transversity PDF


- E.g. $h_1(x)$ from JLab

[Cocuzza et al (JAM Collaboration), PRD (2024)]



- Effective polarization of quarks that participate in the interaction:

$$\vec{b}_{\perp}^{\langle q \rangle} = \alpha \vec{b}_{\perp}^p, \quad \alpha \equiv \frac{\sum_q e_q^2 h_{1,q}(x)}{\sum_q e_q^2 f_q(x)}$$





Parton-level vs Hadron-level Kinematics


- Controlling/Selecting quark transverse polarization from DIS kinematics



Control the quantum state production



$$\vec{b}_{\perp}^{\langle q \rangle} = \alpha \vec{b}_{\perp}^p, \quad \alpha \equiv \frac{\sum_q e_q^2 h_{1,q}(x)}{\sum_q e_q^2 f_q(x)}$$



Summary

- In terms of quantum information, transverse polarization of initial states usually makes a qualitative effect compared to longitudinal polarization. [\[see also Hao Zhang's talk\]](#)
- At EIC, there are plenty of ways to select/control initial states.
 - ▶ e.g. with the transversity PDF, we show the opportunity to control the state production from the initial state polarization.
 - ▶ Beyond leading order, $b_{\perp}^q$ can arise from unpolarized protons in TMD factorization
[\[EIC white paper, arXiv:1212.1701\]](#)
- Spin of quark can be measured. [\[see also Bin Yan's talk\]](#) Looking to the future, direct spin measurements of stable particles at colliders would make more information available, and help understanding QCD or NP?

Quantum entanglement and magic

- Entanglement asks whether the state can be written in a separable form

$$|\psi^{AB}\rangle = |\psi^A\rangle \otimes |\psi^B\rangle \quad \rho^{AB} = \sum_i p_i \rho_i^A \otimes \rho_i^B$$

- ▶ Concurrence, a quantitative measure of entanglement:

$$\mathcal{C}(\rho) = \max(0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4) > 0 \quad \mathcal{C}(\rho) = 0 \text{ (separable)}$$
$$\lambda_i \text{ are eigenvalues of } \sqrt{\sqrt{\rho}\tilde{\rho}\sqrt{\rho}} \quad = 1 \text{ (maximally entangled)}$$

[W. Wootters, PRL (1998)]

$$\tilde{\rho} = (\sigma_2 \otimes \sigma_2) \rho^* (\sigma_2 \otimes \sigma_2)$$

- Magic asks whether how different the state is from stabilizer states generated by Clifford gates


$$\mathcal{P}_n = P_1 \otimes P_2 \otimes \dots \otimes P_n, \quad P_i \in \{\mathbb{1}, \sigma_1, \sigma_2, \sigma_3\},$$

- ▶ Stabilizer Rényi entropy, a quantitative measure of magic

$$M_k(\rho) = -\frac{1}{1-k} \log_2 \left(\frac{\sum_{P \in \mathcal{P}_n} (\text{Tr}[P\rho])^{2k}}{\sum_{P \in \mathcal{P}_n} (\text{Tr}[P\rho])^2} \right) \quad [\text{Leone, Oliviero, and Hamma, PRL (2022)}]$$

Many ways to infer the quark spin

- Depending on different theory inputs:
 - ▶ From production kinematics [KC, T. Han and M. Low, PLB (2025)]
 - ▶ From Λ hadron spin ($q \rightarrow \Lambda_q + X, \Lambda_q \rightarrow \dots$) [Afik, Kats, de Nova, Soffer and Uzan, PRD (2025)]
[Lin, Liu, Shao, and Wei, JHEP (2025)]
(some spin fraction of q is carried by Λ_q , nonperturbative theory inputs)
 - ▶ From interference fragmentation functions ($q \rightarrow \pi^+ \pi^- + X$) [KC and Yan, PRL (2025)]



Backup: Elastic scattering

- Elastic, form factors $J_{e,\alpha\alpha'}^\mu = \bar{u}_{\alpha'}(p'_e)\gamma^\mu u_\alpha(p_e), \quad J_{p,\beta\beta'}^\mu = \bar{u}_{\beta'}(p'_p)\Gamma^\mu u_\beta(p_p),$

$$\Gamma^\mu = \gamma^\mu F_1(q^2) + \frac{i\sigma^{\mu\nu}q_\nu}{2m_p} F_2(q^2), \quad q = p'_p - p_p, \quad \sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu].$$

$$\begin{aligned} \mathcal{T} &= \begin{pmatrix} \mathcal{M}_{\uparrow\uparrow\uparrow\uparrow} & \mathcal{M}_{\uparrow\uparrow\uparrow\downarrow} & 0 & 0 \\ \mathcal{M}_{\uparrow\uparrow\downarrow\uparrow} & \mathcal{M}_{\uparrow\uparrow\downarrow\downarrow} & 0 & 0 \\ 0 & 0 & \mathcal{M}_{\downarrow\downarrow\uparrow\uparrow} & \mathcal{M}_{\downarrow\downarrow\uparrow\downarrow} \\ 0 & 0 & \mathcal{M}_{\downarrow\downarrow\downarrow\uparrow} & \mathcal{M}_{\downarrow\downarrow\downarrow\downarrow} \end{pmatrix} \\ &= 2e^2 \frac{E_e E_p}{t} F_1(q^2) \begin{pmatrix} e^{i\phi}(1+\beta)(1+c_\theta) & s_\theta \sqrt{1-\beta^2} & 0 & 0 \\ -e^{i\phi} s_\theta \sqrt{1-\beta^2} & 1+3\beta+(1-\beta)c_\theta & 0 & 0 \\ 0 & 0 & 1+3\beta+(1-\beta)c_\theta & e^{-i\phi} s_\theta \sqrt{1-\beta^2} \\ 0 & 0 & s_\theta \sqrt{1-\beta^2} & e^{-i\phi}(1+\beta)(1+c_\theta) \end{pmatrix} \\ &\quad + 4e^2 \frac{E_e E_p}{t} F_2(q^2) \begin{pmatrix} 0 & -s_\theta \frac{\beta(\beta+1)}{\sqrt{1-\beta^2}} & 0 & 0 \\ e^{i\phi} s_\theta \frac{\beta(\beta+1)}{\sqrt{1-\beta^2}} & 2\beta(1-c_\theta) & 0 & 0 \\ 0 & 0 & 2\beta(1-c_\theta) & -e^{-i\phi} s_\theta \frac{\beta(\beta+1)}{\sqrt{1-\beta^2}} \\ 0 & 0 & s_\theta \frac{\beta(\beta+1)}{\sqrt{1-\beta^2}} & 0 \end{pmatrix} \end{aligned}$$