



Entanglement dynamics in Deep Inelastic Scattering from real-time quantum simulation

Meijian Li 李枚键

Huazhong University of Science and Technology

Based on arXiv:2610.25350, in collaboration with X. Cao (USTC), Y. Li (USTC), W. Qian (CCNU).




Quantum Entanglement Studies in Nuclear and Particle Physics

October 24-26, 2026, Fudan University, Shanghai, China

Deep Inelastic Scattering


Renewed and growing interests





HERA - H1, Zeus (1992-2007)
Hamburg, Germany

Deep Inelastic Scattering

Renewed and growing interests



HERA - H1, Zeus (1992-2007)
Hamburg, Germany




EIC (2030+)
New York, USA

+ LHeC, EicC, ...


Hadron Production in DIS

- DIS at small x : $\gamma^* \rightarrow q\bar{q}$ dipole $\rightarrow$ nuclear color field $\rightarrow$ hadron
- Multiple scales in one process:
 - Initial splitting: perturbative
 - Hadron formation: nonperturbative
 - Quantum coherence preserved throughout
- Conventional approaches (factorization, MC generators) break quantum coherence
- Goal: first-principles, amplitude-level simulation preserving the full quantum state



$$\gamma^* + p/A \rightarrow J/\psi + X$$

Light Front Quantization



[Figure adapted from Yang Li]

- Bound states are solved from eigenvalue equations

$$P_{QCD}^- |\phi\rangle = P_\phi^- |\phi\rangle \Leftrightarrow \underbrace{(P_{QCD}^- P^+ - \vec{P}_\perp^2)}_{H_{LC} \text{ boost invariant}} |\psi_h(P, j, m_j)\rangle = M_h^2 |\psi_h(P, j, m_j)\rangle$$



- Evolution of a quantum state follows the Schrödinger equation

$$\frac{1}{2} P^- |\psi(x^+)\rangle = i \frac{\partial}{\partial x^+} |\psi(x^+)\rangle$$

LF QCD Hamiltonian *with background*

- **first-principles:** QCD Lagrangian via Legendre transformation,

$$\mathcal{L} = -\frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a + \bar{\psi}(i\gamma^\mu D_\mu - m)\psi, D_\mu = \partial_\mu + ig(A_\mu + \mathcal{A}_\mu) \rightarrow P^-(x^+) = P_{KE}^- + V_{qg} + V_{ggg} + V_{gggg} + W_f + W_g + V_{\mathcal{A}}(x^+)$$



Color Glass Condensate in DIS: $\mathcal{A}_\mu$

strong field, weak coupling,
large occupation numbers

classical fields by Yang-Mills eq.

$$\langle \rho_a(x)\rho_b(y) \rangle = g^2 \tilde{\mu}^2 \delta_{ab} \delta^3(\vec{x} - \vec{y}), \quad (m_g^2 - \nabla_\perp^2) \mathcal{A}_a^-(x^+, \vec{x}_\perp) = \rho_a(x^+, \vec{x}_\perp)$$

$$\text{Saturation scale: } Q_s^2 = C_F (g^2 \tilde{\mu})^2 L_\eta / (2\pi^2)$$

Fock space expansion

- Fock sector expansion of quantum states:

$$|q\rangle_{\text{jet}} = \psi_q|q\rangle + \psi_{qg}|qg\rangle + \psi_{qgg}|qgg\rangle + \dots$$

$$|g\rangle_{\text{jet}} = \psi_g|g\rangle + \psi_{gg}|gg\rangle + \psi_{ggg}|ggg\rangle + \dots$$

$$|\gamma^*\rangle_{\text{phys}} = \psi_\gamma|\gamma^*\rangle + \psi_{q\bar{q}}|q\bar{q}\rangle + \psi_{qqg}|qq\bar{q}\rangle + \dots$$

$$|q(\beta)\rangle = b_\beta^\dagger|0\rangle,$$

$$|\bar{q}(\beta)\rangle = d_\beta^\dagger|0\rangle,$$

$$|g(\beta)\rangle = a_\beta^\dagger|0\rangle.$$

$$|\psi(x^+)\rangle = \sum_{\beta} c_{\beta}(x^+)|\beta\rangle$$

Each single-particle basis state is labeled by five quantum numbers:


$$\beta_l = \{p_l^+, p_l^x, p_l^y, \lambda_l, c_l\}, (l = q, \bar{q})$$

$$|q\rangle: |\beta_q\rangle$$

$$|qg\rangle: |\beta_{qg}\rangle = |\beta_q\rangle \otimes |\beta_g\rangle$$

$$|q\bar{q}\rangle: |\beta_{q\bar{q}}\rangle = |\beta_q\rangle \otimes |\beta_{\bar{q}}\rangle$$

...



Fock space expansion

- Fock sector expansion of quantum states:

$$|q\rangle_{\text{jet}} = \psi_q |q\rangle + \psi_{qg} |qg\rangle + \psi_{qgg} |qgg\rangle + \dots$$

$$|q(\beta)\rangle = b_\beta^\dagger |0\rangle,$$

$$|g\rangle_{\text{jet}} = \psi_g |g\rangle + \psi_{gg} |gg\rangle + \psi_{ggg} |ggg\rangle + \dots$$

$$|\bar{q}(\beta)\rangle = d_\beta^\dagger |0\rangle,$$

$$|g(\beta)\rangle = a_\beta^\dagger |0\rangle.$$

$$|\gamma^*\rangle_{\text{phys}} = \psi_\gamma |\gamma^*\rangle + \psi_{q\bar{q}} |q\bar{q}\rangle + \psi_{qqg} |qqg\rangle + \dots$$

$$|\psi(x^+)\rangle = \sum_{\beta} c_{\beta}(x^+) |\beta\rangle$$

Each single-particle basis state is labeled by five quantum numbers:


$$\beta_l = \{p_l^+, p_l^x, p_l^y, \lambda_l, c_l\}, (l = q, \bar{q})$$

$$|q\rangle: |\beta_q\rangle$$


$$|qg\rangle: |\beta_{qg}\rangle = |\beta_q\rangle \otimes |\beta_g\rangle$$

$$|q\bar{q}\rangle: |\beta_{q\bar{q}}\rangle = |\beta_q\rangle \otimes |\beta_{\bar{q}}\rangle$$

...



Quantum implementation: Encoding



Time evolution of quantum states

- **Non-perturbative:** time evolution operator into a product of many sequential timesteps


$$|\psi(x^+)\rangle = \mathcal{T}_+ e^{-\frac{i}{2} \int_0^{x^+} dz^+ P^-(z^+)} |\psi(0)\rangle$$

$\equiv U(x^+; 0)$


Trotterization

$$\prod_{k=1}^n \mathcal{T}_+ \exp \left[-\frac{i}{2} \int_{x_{k-1}^+}^{x_k^+} dz^+ P^-(z^+) \right] |\psi(0)\rangle$$

in which $\delta x^+ = x^+/n$, $x_k^+ = k\delta x^+$ ($k = 0, 1, 2, \dots, n$).



automatically **going beyond eikonal**
and include **quantum interference**
at each timestep:



Time evolution of quantum states

- **Non-perturbative:** time evolution operator into a product of many sequential timesteps


$$|\psi(x^+)\rangle = \mathcal{T}_+ e^{-\frac{i}{2} \int_0^{x^+} dz^+ P^-(z^+)} |\psi(0)\rangle$$

$\equiv U(x^+; 0)$

Trotterization


$$\prod_{k=1}^n \mathcal{T}_+ \exp \left[-\frac{i}{2} \int_{x_{k-1}^+}^{x_k^+} dz^+ P^-(z^+) \right] |\psi(0)\rangle$$

in which $\delta x^+ = x^+/n$, $x_k^+ = k\delta x^+$ ($k = 0, 1, 2, \dots, n$).




automatically **going beyond eikonal**
and include **quantum interference**
at each timestep:

Quantum implementation: trotterization




Extraction of physical quantities

- Evaluation of physical quantities is **directly** from the quantum state : $O(x^+) = \langle \psi(x^+) | \hat{O} | \psi(x^+) \rangle$



Extraction of physical quantities

- Evaluation of physical quantities is **directly** from the quantum state : $O(x^+) = \langle \psi(x^+) | \hat{O} | \psi(x^+) \rangle$



Quantum implementation: measurement


Quantum state Amplitude

$$\begin{pmatrix} c_1 \\ \vdots \\ c_i \\ \vdots \\ c_n \end{pmatrix}$$


Classical state

$$P_i = |c_i|^2$$


$$\begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$$



Progress: Real-time Jet classical quantum Simulations



Progress: Real-time Jet classical quantum Simulations




Initial States

Single-momentum dipole

- $\vec{p}_{q,\perp} = \vec{p}_{\bar{q},\perp} = \vec{0}_\perp$, color singlet
- Benchmark against analytical expectation

$$|\psi_{q\bar{q}}\rangle = \sum_{\beta} \sum_{\beta'} c_{\beta\beta'} |q(\beta)\bar{q}(\beta')\rangle$$


$$c_{\beta\beta'} = \begin{cases} \frac{1}{\sqrt{N_c}}, & \text{for } k^i = \bar{k}^i = 0, z = 1/2, k^+ = \bar{k}^+ = K/2, c = \bar{c} \\ 0, & \text{otherwise} \end{cases}$$



Virtual photon

- $q\bar{q}$ component of a photon state
- Physical DIS probe

$$|\gamma^*\rangle_{phys} = |\gamma^*\rangle + |q\bar{q}\rangle + |q\bar{q}g\rangle + \dots$$



$$\psi_{s\bar{s}/\gamma}^{(m_j=0)}(\vec{k}_\perp, x, Q) = Q_f e \sqrt{N_c} \sqrt{x(1-x)} \frac{2Qx(1-x)}{\epsilon^2 + k_\perp^2} \delta_{s,\bar{s}},$$

Initial States and nuclear target


Single-momentum dipole

- $\vec{p}_{q,\perp} = \vec{p}_{\bar{q},\perp} = \vec{0}_\perp$, color singlet
- Benchmark against analytical expectation

Virtual photon

- $q\bar{q}$ component of a photon state
- Physical DIS probe

$$|\gamma^*\rangle_{phys} = |\gamma^*\rangle + |q\bar{q}\rangle + |q\bar{q}g\rangle + \dots$$




$$\text{Saturation scale: } Q_s^2 = C_F(g^2\tilde{\mu})^2 L_\eta / (2\pi^2)$$


$$L_{\eta,A} = A^{1/3} L_{\eta,p}$$

Momentum Broadening

Single-momentum dipole: q




$\bar{q}$




- Classical & quantum-circuit simulations agree
- Early times: linear growth, follows discrete analytical prediction
- Late times: saturates at $\langle p_{\perp}^2 \rangle / d_p^2 \approx 2N_{\perp}^2/3$
 - Finite-basis effect: wraps around periodic boundary

Evolution of momentum distribution

Single-momentum dipole




- Quark momentum broadens over time



Evolution of momentum distribution


Single-momentum dipole



- quark momentum broadens over time


$$\vec{k}_\perp = (\vec{p}_{q,\perp} - \vec{p}_{\bar{q},\perp})/2$$

- relative momentum broadens



Evolution of momentum distribution

Single-momentum dipole




- quark momentum broadens over time

$$\vec{k}_\perp = (\vec{p}_{q,\perp} - \vec{p}_{\bar{q},\perp})/2$$

- relative momentum broadens


$$\vec{P}_\perp = \vec{p}_{q,\perp} + \vec{p}_{\bar{q},\perp}$$

- C.M. momentum broadens faster
 - Peak at $\vec{P}_\perp = \vec{0}_\perp$: coincident component, color transparent




Evolution of momentum distribution

Virtual photon state




- quark momentum broadens over time



Evolution of momentum distribution


Virtual photon state



- quark momentum broadens over time


$$\vec{k}_\perp = (1 - x)\vec{p}_{q,\perp} - x\vec{p}_{\bar{q},\perp}, x = p_q^+ / P^+$$

- relative momentum broadens



Evolution of momentum distribution

Virtual photon state




- quark momentum broadens over time

$$\vec{k}_\perp = (1 - x)\vec{p}_{q,\perp} - x\vec{p}_{\bar{q},\perp}, x = p_q^+ / P^+$$

- relative momentum broadens

$$\vec{P}_\perp = \vec{p}_{q,\perp} + \vec{p}_{\bar{q},\perp}$$

- C.M. momentum barely broadens
 - $r_\perp \approx 0.73 \text{ GeV}^{-1} \ll a_\perp \approx 3 \text{ GeV}^{-1}$, strong correlation between q and $\bar{q}$



Cross Section

Cross section is defined as

$$\frac{d\sigma}{d^2b} = \sum_{\phi_f} |M(\phi_f; \psi_{in})|^2$$

finite-time S-matrix


$$\begin{aligned} M(\phi_f; \psi_{in}) &= \langle \phi_f | S | \psi_{in} \rangle - \langle \phi_f | \psi_{in} \rangle \\ &= \langle \phi_f | \psi_{out} \rangle - \langle \phi_f | \psi_{in} \rangle \end{aligned}$$

Cross Section

Cross section is defined as

$$\frac{d\sigma}{d^2b} = \sum_{\phi_f} |M(\phi_f; \psi_{in})|^2$$

finite-time S-matrix

$$\begin{aligned} M(\phi_f; \psi_{in}) &= \langle \phi_f | S | \psi_{in} \rangle - \langle \phi_f | \psi_{in} \rangle \\ &= \langle \phi_f | \psi_{out} \rangle - \langle \phi_f | \psi_{in} \rangle \end{aligned}$$

- **total cross section**

$$\frac{d\sigma_{\text{tot}}}{d^2b} = \left\langle \sum_{\phi_f} |M(\phi_f; \psi_{in})|^2 \right\rangle_{\text{event}}$$

- **elastic cross section**

$$\frac{d\sigma_{\text{el}}}{d^2b} = \sum_{\phi_f} |\langle M(\phi_f; \psi_{in}) \rangle_{\text{event}}|^2$$

event average = medium fluctuations from nuclear target

Cross Section


- total cross section

$$\frac{d\sigma_{\text{tot}}}{d^2b} = \langle \sum_{\phi_f} |M(\phi_f; \psi_{in})|^2 \rangle_{\text{event}}$$



$$\frac{d\sigma_{\text{tot}}}{d^2b} = 2 (1 - \langle \text{Re} \langle \psi_{in} | \psi_{out} \rangle \rangle_{\text{event}})$$

- Quantum implementation via Hadamard Tests



- elastic cross section

$$\frac{d\sigma_{\text{el}}}{d^2b} = \sum_{\phi_f} |\langle M(\phi_f; \psi_{in}) \rangle_{\text{event}}|^2$$





$$\frac{d\sigma_{\text{el}}}{d^2b} = |\langle \text{Re} \langle \psi_{in} | \psi_{out} \rangle - 1 \rangle_{\text{event}}|^2 + |\langle \text{Im} \langle \psi_{in} | \psi_{out} \rangle \rangle_{\text{event}}|^2$$

$$\text{Re} \langle \psi_{in} | \psi_{out} \rangle = P(0) - P(1)$$

$$\text{Im} \langle \psi_{in} | \psi_{out} \rangle = P(0) - P(1)$$

Cross Section

Single-momentum dipole





- Simulation results agree with analytical expectation
- Approaches black-disk limit ($d\sigma_{tot}/d^2b \rightarrow 2$, $d\sigma_{el}/d^2b \rightarrow 1$) at later time

$$\left. \frac{d\sigma_{q\bar{q},tot}}{d^2b} \right|_{p^+=\infty} = 2 \left\{ 1 - \exp \left[-C_F \frac{g^4 \tilde{\mu}^2 L_\eta}{4\pi m_q^2} [1 - m_g |\vec{x}_\perp - \vec{y}_\perp| K_1(m_g |\vec{x}_\perp - \vec{y}_\perp|)] \right] \right\}$$

$$\left. \frac{d\sigma_{q\bar{q},el}}{d^2b} \right|_{p^+=\infty} = \left\{ 1 - \exp \left[-C_F \frac{g^4 \tilde{\mu}^2 L_\eta}{4\pi m_g^2} [1 - m_g |\vec{x}_\perp - \vec{y}_\perp| K_1(m_g |\vec{x}_\perp - \vec{y}_\perp|)] \right] \right\}^2$$

Cross Section

Virtual photon state




- Simulation results agree with analytical expectation
- Approaches black-disk limit ($d\sigma_{tot}/d^2b \rightarrow 2 d\sigma_{el}/d^2b$) at later time for all Q^2
- Slower saturation compared to single-momentum dipole
- Non-monotonic Q^2 dependence: amplitude growth v.s. smaller dipole size

Hadron Production

The probability to produce a hadron via projection

$$\text{Prob}(h) = \sum_{\vec{P}_\perp} |\langle \psi(x^+) | \psi_h(\vec{P}_\perp) \rangle|^2$$




Hadron Production

The probability to produce a hadron via projection

$$\text{Prob}(h) = \sum_{\vec{P}_\perp} |\langle \psi(x^+) | \psi_h(\vec{P}_\perp) \rangle|^2$$

The associated nuclear modification factor: $R_A = \frac{\text{Prob}_A(J/\psi)}{\text{Prob}_p(J/\psi)}$






Hadron Production

The probability to produce a hadron via projection



$$\text{Prob}(h) = \sum_{\vec{P}_\perp} |\langle \psi(x^+) | \psi_h(\vec{P}_\perp) \rangle|^2$$

The associated nuclear modification factor: $R_A = \frac{\text{Prob}_A(J/\psi)}{\text{Prob}_p(J/\psi)}$

- Quantum implementation via SWAP test or inversion test





J/ψ Production



- $\text{Prob}(J/\psi)$ maximal at early x^+ , decreases as medium scrambles the state
- Non-monotonic in Q^2 :
 - small Q^2 : photon amplitude grows
 - large Q^2 : wavefunction broadens

J/ψ Production



- Production dominated by $P_{\perp} \approx 0$ at both Q^2
 - color transparency, compact singlet barely kicked
- $R_A \approx 1$ at lowest and larger $P_{\perp}$
- Suppression at $P_{\perp} \sim 0.25\text{--}0.5 \text{ GeV}$

Entanglement Entropy


- The von Neumann (vN) entropy of the reduced density matrix obtained by tracing out the antiquark degrees of freedom

$$S_{\text{vN}}(x^+) = -\text{Tr}[\rho(x^+)\log_2\rho(x^+)]$$

- Normalize $q\bar{q}$ state to unity before tracing
- Trace out antiquark degrees of freedom

$$\rho(x^+) = \text{Tr}_{\bar{q}}(|\psi(x^+)\rangle\langle\psi(x^+)|)$$

- Event average $\langle\rho(x^+)\rangle_{event}$



Entanglement Entropy

- The von Neumann (vN) entropy of the reduced density matrix obtained by tracing out the antiquark degrees of freedom

$$S_{\text{vN}}(x^+) = -\text{Tr}[\rho(x^+)\log_2\rho(x^+)]$$


- Normalize $q\bar{q}$ state to unity before tracing
- Trace out antiquark degrees of freedom

$$\rho(x^+) = \text{Tr}_{\bar{q}}(|\psi(x^+)\rangle\langle\psi(x^+)|)$$

- Event average $\langle\rho(x^+)\rangle_{event}$

- On the discrete momentum basis representation


- q Hilbert space: $K \times N_c (2N_\perp)^2$ modes
- p_q^+ conserved $\rightarrow$ longitudinal resolution K does not enter
- Bounds: $0 \leq S_{\text{vN}} \leq \log_2[N_c (2N_\perp)^2]$



S_{vN} measures entanglement between quark and antiquark in color $\times$ momentum space, induced by color rotations and momentum kicks by medium.

Entanglement Entropy


Single-momentum dipole



- $S_{vN}(0) = \log_2 N_c \approx 1.6$
 - pure color entanglement of the singlet dipole
- Rapid growth due to medium interaction
- Asymptotic value: $\log_2 [N_c (2N_\perp)^2] \approx 7.6$, quark fully randomized


Entanglement Entropy

Single-momentum dipole



- $S_{vN}(0) = \log_2 N_c \approx 1.6$
 - pure color entanglement of the singlet dipole
- Rapid growth due to medium interaction
- Asymptotic value: $\log_2 [N_c (2N_\perp)^2] \approx 7.6$, quark fully randomized


Virtual photon state



- Initial entropy already large, $S_{vN}(0) \approx 7.5$
 - Photon WF superposes many color $\times$ momentum modes
- Growth is modest but reaches same asymptotic value

Entanglement Entropy

Connection to observables



Entanglement entropy provides an observable-independent diagnostic of how strongly the medium scrambles the dipole system

Summary and Outlook

- ✓ We developed and applied a **non-perturbative** computational approach to simulate the **real-time** evolution of dipole-to-hadron production in DIS
 1. Quantum simulation framework developed for in-medium dipole evolution
 2. Cross sections agree with analytical expectations
 3. J/ψ production and R_A
 4. Entanglement entropy as an observable-independent diagnostic of medium effect

Next:

- Non-eikonal, $P^+ < \infty$
- Inclusion of higher Fock sector contribution, $|\gamma^*\rangle_{phys} = |\gamma^*\rangle + |q\bar{q}\rangle + |q\bar{q}g\rangle + \dots$
- Photon state preparation on quantum circuit

Thank you!